

Comment on “Ising Spin Glasses in a Magnetic Field”

In a very interesting paper [1] Houdayer and Martin analyze the $T = 0$ $3d$ EA spin glass with a magnetic field B . By using a new, powerful method, they determine an effective critical field B_c (given by the crossings of their function r , the rate at which ground states change) as a function of the lattice size L . They use their results to deduce that the model is behaving like in the droplet approach and not like the mean-field theory, that implies the survival of a phase transition for finite field B . We show here, by using some unpublished data [2], that the very interesting method and numerical results of [1] are completely compatible with the behavior implied by the Replica Symmetry Breaking (RSB) theory.

In [3] we have introduced a dynamical approach that has allowed us to determine the value of the minimal allowed overlap, q_{min} , for the $4d$ EA model in a non-zero magnetic field. By establishing that this value is different from q_{ave} , the average overlap measured on medium size samples, we have shown that also in $4d$ the phase transition to a RSB phase survives after applying a magnetic field. We use here analogous data for the $3d$ case [2], that also clearly show the survival of the Almeida-Thouless line, to interpret the result of [1].




FIG. 1. $q(t_A)$ versus t_A at $T = 0.5$, $B = 0.4$.

In our approach we use a large lattice (for example $L = 32$), and we select an annealing schedule to run a Metropolis dynamics. We select a set of decreasing values of the temperature $\{T_n\}$, starting from high $T \gg T_c$, and we spend $n \cdot 2^k$ updating steps at each T_n value. We compute $q(t_A)$, where $t_A \equiv 2^k$, for increasing k values. The limit for $t_A \rightarrow \infty$ of $q(t_A)$ is just q_{min} . The value of q_{ave} is computed with usual equilibrium tempering simulations on smaller lattices. We show in figure (1) what happens in $3d$ at $B = 0.4$ and $T = 0.5$, where RSB is clear: the fit of the data is very precise, and gives a value of q_{min} that clearly differs from q_{ave} . Ref. [1] deals with $T = 0$, so in figure (2) we extrapolate $q_{min}(T)$ at $B = 0.4$ down to $T = 0$. A simple power fit is very good, giving

$q_{min}(T = 0) = 0.74$. In terms of the percentage of different spins x of [1] this means that at $B = 0.4$ we find $x \simeq 0.13$, corresponding to a jump with overlap 0.74: in the conditions of [1] (where $x = 0.15$) one expects as $L \rightarrow \infty$ a crossing of the r curves close to $B_c = 0.4$.




FIG. 2. $q_{min}(T)$ and best power fit.

This is well compatible with the data of [1]: their crossing point moves from $B \simeq 2.1$ to $B \simeq 1.2$ when L goes from 3 to 8: in these conditions it is difficult to discriminate between the droplet $B_c = 0$ and the $B_c \simeq 0.4$ that follows from RSB. RSB predicts that $r \simeq L^{\frac{3}{2}}$ at small B , which is not far from the observed behavior. A main feature of short range models is that one needs very large L values to determine the asymptotic value of q_{min} : the value of q_{min} becomes sizably different from zero already for a small B ($q_{min} \approx B^{0.3}$ in $4d$ against $q_{min} \approx B^{\frac{3}{2}}$ in $d = \infty$). This implies that the width at $L = \infty$ of $P(q)$ becomes much smaller as soon as B is switched on, and finite volume effects become much more important. This effect is exacerbated by the presence of the very large, low q tail of $P(q)$ at L not too large (see [4] for the analogous $4d$ case), that disappears very slowly for increasing L . The new method of [1] is very promising, and going to larger lattices will probably allow one further verification of RSB for finite d spin glasses.

These simulations have been run at ZIB-Berlin. We thank H. Stueben for his crucial and extensive help: without him this work would have been impossible. We acknowledge W. Janke and D. Johnston contribution in setting up this project.

E. Marinari, G. Parisi and F. Zuliani

-
- [1] J. Houdayer and O. C. Martin, cond-mat/9811419.
 - [2] E. Marinari, G. Parisi and F. Zuliani, in preparation.
 - [3] E. Marinari, G. Parisi and F. Zuliani, J. Phys. A **31** (1998) 1181, cond-mat/9703253.
 - [4] E. Marinari, C. Naitza and F. Zuliani, J. Phys. A **31** (1998) 6355, cond-mat/9802224.